



Mathematics and CS Seminar

Hyperbolic branching Brownian motion

Wolfgang Woess

TU Graz

Host: Laszlo Erdös, Jan Maas

Euclidean branching Brownian motion (BBM) has been intensively studied during many decades by renowned researchers. BBM on hyperbolic space has received less attention. A profound study of Lalley and Sellke (1997) provided insight on the recurrent, resp. transient regimes of BBM on the Poincare' disk. In particular, they determined the Hausdorff dimension of the limit set on the boundary circle in dependence on the fission rate of the branching particles. In the present notes, further features are exhibited. The rates of the maximal and minimal hyperbolic distances to the starting point are determined, as well as refined asymptotic estimates in the transient regime. The other main issues studied here concern the behaviour of the empiricial distributions of the branching population, as time goes to infinity, and their convergence to an infinitely supported random limit probability measure on the boundary.

Monday, December 16, 2024 05:00pm - 06:00pm

Central Bldg / O1 / Mondi 2a (I01.O1.008)



This invitation is valid as a ticket for the ISTA Shuttle from and to Heiligenstadt Station. Please find a schedule of the ISTA Shuttle on our webpage: https://ista.ac.at/en/campus/how-to-get-here/ The ISTA Shuttle bus is marked ISTA Shuttle (#142) and has the Institute Logo printed on the side.

www.ista.ac.at | Institute of Science and Technology Austria | Am Campus 1 | 3400 Klosterneuburg