



Seminar/Talk

S-dual of Hamiltonian G spaces and relative Langlands duality

Hiraku Nakajima

IPMU

Host: Tamas Hausel

The S-dual (G, M) of the pair (G, M) of a smooth affine algebraic symplectic manifold M with hamiltonian action of a complex reductive group G was introduced implicitly in [arXiv:1706.02112] and explicitly in [arXiv:1807.09038] under the cotangent type assumption. The definition was a modification of the definition of Coulomb branches of gauge theories in [arXiv:1601.03586]. It was motivated by the S-duality of boundary conditions of 4-dimensional $N=4$ super Yang-Mills theory, studied by Gaiotto and Witten [arXiv:0807.3720]. It is also relevant to the relative Langlands duality proposed by Ben-Zvi, Sakellaridis and Venkatesh. In this article, we review the definition and properties of S-dual.

Thursday, October 3, 2024 01:00pm - 03:00pm

Office Bldg West / Ground floor / Heinzl Seminar Room (I21.EG.101)



This invitation is valid as a ticket for the ISTA Shuttle from and to Heiligenstadt Station. Please find a schedule of the ISTA Shuttle on our webpage: <https://ista.ac.at/en/campus/how-to-get-here/> The ISTA Shuttle bus is marked ISTA Shuttle (#142) and has the Institute Logo printed on the side.