



Seminar/Talk

Cobordism maps in knot Floer homology

Andras Juhasz

U of Oxford

Host: Tamas Hausel

An n -manifold is a topological space that locally looks like n -dimensional coordinate space. Surprisingly, the most difficult dimensions to understand are 3 and 4. Low-dimensional topology is an important area of mathematics that studies manifolds in exactly these dimensions. Knots play a central role in low-dimensional topology as they can be used to construct all 3- and 4-manifolds, and they also appear in physics, biology, and chemistry. Knot Floer homology is a powerful, computable, and geometrically rich invariant of knots defined by Ozsvath-Szabo and Rasmussen in 2002. Some of its properties are best understood via sutured Floer homology, a generalization to 3-manifolds with boundary that I developed. It is a fundamental question of low-dimensional topology to understand the surfaces a knot can bound in the 4-ball. In this talk, I will explain how a knot cobordism (a surface in 4-space connecting two knots) induces a functorial map on knot Floer homology via sutured Floer homology, and discuss some properties and applications.

Tuesday, March 13, 2018 11:00am - 12:00pm

Big Seminar room Ground floor / Office Bldg West (I21.EG.101)



This invitation is valid as a ticket for the ISTA Shuttle from and to Heiligenstadt Station.

Please find a schedule of the ISTA Shuttle on our webpage:

<https://ista.ac.at/en/campus/how-to-get-here/> The ISTA Shuttle bus is marked ISTA Shuttle (#142) and has the Institute Logo printed on the side.