



Mathematics and CS Seminar

Percolation Phase Transition via the Gaussian Free Field

Aran Raoufi

ETH Zürich

Host: M. Beiglböck, N. Berestycki, L. Erdős, J. Maas

Let G be a bounded-degree infinite graph, and p_c be the critical parameter of bond percolation on G . That is p_c is the infimum of values of p that you have an infinite cluster almost surely. In this talk, we prove that if the isoperimetric dimension of G is higher than 4, then $p_c(G) < 1$. The theorem settles affirmatively two conjectures of Benjamini and Schramm. Notably, if G is a transitive graph with super-linear growth, then $p_c(G) < 1$. In particular, it implies that if G is a Cayley graph of a finitely generated group without a finite index cyclic subgroup, then $p_c(G) < 1$. The proof of the theorem starts with the existence of an infinite cluster for percolation in a certain in-homogeneous random environment governed by the Gaussian free field. Then, by the help of a multiscale decomposition of GFF, we relate the existence of an infinite cluster in percolation in the random environment to that of percolation with a fixed parameter $p < 1$. This talk is based on a joint work with H. Duminil-Copin, S. Goswami, F. Severo, and A. Yadin.

Tuesday, May 21, 2019 05:30pm - 06:30pm

Big Seminar room Ground floor / Office Bldg West (I21.EG.101)



This invitation is valid as a ticket for the ISTA Shuttle from and to Heiligenstadt Station.

Please find a schedule of the ISTA Shuttle on our webpage:

<https://ista.ac.at/en/campus/how-to-get-here/> The ISTA Shuttle bus is marked ISTA Shuttle (#142) and has the Institute Logo printed on the side.