



Seminar/Talk

GeomTop Seminar: "Chromatic number of Kneser hypergraphs and a conjecture of Frick"

Amir Jafari

Host: Uli Wagner

This will be a report of a recent joint work with Soheil Azarpendar. For integers n, k, r where $n > kr - 1$ The Kneser hypergraph $KG^r(n, k)$ was defined by Lovasz, Alon and Frankl as the r -uniform hypergraph with all k -subsets of $\{1, \dots, n\}$ as vertices and its hyperedges are all r -subsets $\{A_1, \dots, A_r\}$ of vertices that are pairwise disjoint. It was proved by them using topological methods that its chromatic number is the ceiling of $(n - r(k - 1)) / (r - 1)$. Ziegler conjectured that if we take the induced sub hypergraph whose vertices are all r -stable (i.e subsets that for any two distinct elements i and j in them $r - 1 < |i - j| < 2$). In this talk we prove a weaker version of this conjecture, due to Frick et al, that states if $\{P_1, \dots, P_t\}$ is a partition of $\{1, \dots, n\}$ where the size of each P_i is at most r and we take the induced sub hypergraph of $KG^r(n, k)$ whose vertices are those k -subsets that have at most one element from each P_i then we still get the same chromatic number. Our proof for this conjecture is combinatorial and uses \mathbb{Z}_p Tucker lemma. If time permits some topological methods related to similar problems will be explained.

Wednesday, March 4, 2020 01:00pm - 02:15pm

Mondi Seminar Room 3, Central Building



This invitation is valid as a ticket for the ISTA Shuttle from and to Heiligenstadt Station. Please find a schedule of the ISTA Shuttle on our webpage: <https://ista.ac.at/en/campus/how-to-get-here/> The ISTA Shuttle bus is marked ISTA Shuttle (#142) and has the Institute Logo printed on the side.