



Seminar/Talk

An arithmetic count of rational plane curves

Kirsten Wickelgren

Duke University, North Carolina

Host: Timothy Browning

There are finitely many degree d rational plane curves passing through $3d-1$ points, and over the complex numbers, this number is independent of (generically) chosen points. For example, there are 12 degree 3 rational curves through 8 points, one conic passing through 5, and one line passing through 2. Over the real numbers, one can obtain a fixed number by weighting real rational curves by their Welschinger invariant, and work of Solomon identifies this invariant with a local degree. It is a feature of A1-homotopy theory that analogous real and complex results can indicate the presence of a common generalization, valid over a general field. We develop and compute an A1-degree, following Morel, of the evaluation map on Kontsevich moduli space to obtain an arithmetic count of rational plane curves, which is valid for any field k of characteristic not 2 or 3. This shows independence of the count on the choice of generically chosen points with fixed residue fields, strengthening a count of Marc Levine. This is joint work with Jesse Kass, Marc Levine, and Jake Solomon.

Thursday, January 14, 2021 03:00pm - 04:00pm

<https://mathseminars.org/seminar/AGNTISTA>



This invitation is valid as a ticket for the ISTA Shuttle from and to Heiligenstadt Station.

Please find a schedule of the ISTA Shuttle on our webpage:

<https://ista.ac.at/en/campus/how-to-get-here/> The ISTA Shuttle bus is marked ISTA Shuttle (#142) and has the Institute Logo printed on the side.